

Speaker's Profile

-Yukio Yuzawa-

Speaker's Name	Yukio Yuzawa	Country	Japan	
Organization	Fujita Health Univ.	Department	Department of Nephrology	
Education	<p>1987: Ph.D., Nephrology, Nagoya Univ. School of Medicine, Internal Medicine III, Nagoya, Japan (Advisor: Prof. Nobuo Sakamoto, M.D., Ph.D.)</p> <p>1975-1981: M.D., Nagoya Univ. School of Medicine, Nagoya, Japan</p>			
Experience	<p>2010-Present: Professor, Department of Nephrology, Fujita Health Univ.</p> <p>2007-2010: Chief of Nephrology Unit, Nagoya Univ. Hospital</p> <p>2002-2010: Associate Professor, Department of Nephrology, Nagoya Univ. Graduate School of Medicine</p> <p>1990-2002: Instructor, Department of Nephrology, Nagoya Univ. Graduate School of Medicine</p> <p>1987-1990: Visiting Associate Professor, Department of Pathology, State Univ. of New York at Buffalo (Prof. Guiseppe Andres, Prof. Jan Brentjens)</p> <p>1986-1987: Instructor, Internal Medicine III, Nagoya Univ. School of Medicine</p> <p>1983-1986: Staff Physician, Nephrology, Nagoya Univ. Hospital</p> <p>1982-1983: Fellow, Internal Medicine, Nagoya 1st Red-Cross Hospital, Aichi, Japan</p> <p>1981-1982: Resident, Internal Medicine, Nagoya 1st Red-Cross Hospital, Aichi, Japan</p>			
Main Specific Publication	<ul style="list-style-type: none"> - Ishii H, Kumada Y, Toriyama T, Aoyama T, Takahashi H, Tanaka M, Kamoi D, Kawamura Y, Yamada S, Hayashi M, Yasuda Y, Yuzawa Y, Maruyama S, Matsuo S, Matsubara T, Murohara T. : Effects of oral cilostazol 100 mg BID on long-term patency after percutaneous transluminal angioplasty in patients with femoropopliteal disease undergoing hemodialysis: A retrospective chart review in Japanese patients. Clin Ther ;32(1):24-33, 2010 - Okada R, Yuzawa Y, Kawamura T, Hamajima N, Watanabe Y, Matsuo S. Incidence of fever of unknown origin and subsequent antitubercular medications in hemodialysis patients: a two-year prospective study. Ren Fail. 31(10):863-8. 2009 - Kurata K, Maruyama S, Kato S, Sato W, Yamamoto J, Ozaki T, Nitta A, Nabeshima T, Morita Y, Mizuno M, Ito Y, Yuzawa Y, Matsuo S. Tissue-type plasminogen activator deficiency attenuates peritoneal fibrosis in mice. Am J Physiol Renal Physiol. 297(6):F1510-7. 2009 - Ishii H, Toriyama T, Aoyama T, Takahashi H, Amano T, Hayashi M, Tanaka M, Kawamura Y, Yasuda Y, Yuzawa Y, Maruyama S, Matsuo S, Matsubara T, Murohara T. Prognostic values of C-reactive protein levels on clinical outcome after implantation of sirolimus-eluting stents in patients on hemodialysis. Circ Cardiovasc Interv. 2(6):513-8. 2009 			
Summary	<p>Recent my research fields are mainly on 1) the patho-physiological mechanisms of progressive renal injuries including GN, diabetic nephropathy and lupus nephritis, 2) organ cross-talks in CKD and AKI, 3) immunological dysfunction in ESRD, 4) biomarkers of AKI, diabetic nephropathy and CKD.</p>			

Your profile has been shortened to fit our database system.

For any questions or comments on the edited file, please feel free to contact the APCN2010 Secretariat at apcn@intercom.co.kr